

Mixtilinear Incircles

Author(s): Paul Yiu

Source: *The American Mathematical Monthly*, Vol. 106, No. 10 (Dec., 1999), pp. 952-955

Published by: Mathematical Association of America

Stable URL: <http://www.jstor.org/stable/2589751>

Accessed: 25-07-2016 13:34 UTC

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at

<http://about.jstor.org/terms>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



*Mathematical Association of America* is collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*

and let  $\zeta_1, \dots, \zeta_n$  be the (not necessarily distinct) eigenvalues of  $N$ . We leave the proof of the following assertion as an exercise:

$$|M| = \prod_{r=1}^n p(\zeta_r).$$

---

#### REFERENCES

1. R. Bhatia, R. A. Horn, and F. Kittaneh, Normal approximants to binormal operators, *Linear Alg. Appl.* **147** (1991), 169–179.
2. R. R. Halmos, *Finite-dimensional Vector Spaces*, Springer-Verlag, New York, 1993.
3. R. A. Horn, Solution to Problem 96–11, *SIAM Review* **39** (1997), 518–519.
4. K. Hoffman and R. Kunze, *Linear Algebra*, Prentice-Hall, New Jersey, 1971.
5. D. S. Silver and S. G. Williams, Coloring link diagrams with a continuous palette, *Topology*, to appear.

*Department of Mathematics and Statistics, University of South Alabama, Mobile, AL 36688*  
*silver@mathstat.usouthal.edu, williams@mathstat.usouthal.edu*

---

## Mixtilinear Incircles

---

**Paul Yiu**

---

L. Bankoff [1] has coined the term *mixtilinear incircles* of a triangle for the three circles each tangent to two sides and to the circumcircle internally. Consider a triangle  $ABC$  and its mixtilinear incircle in the angle  $A$ , with center  $K_A$ , and radius  $\rho_A$ . Bankoff has established the fundamental formula

$$r = \rho_A \cdot \cos^2 \frac{\alpha}{2}, \tag{1}$$

where  $r$  is the inradius of the triangle, and  $\alpha$  is the magnitude of the angle at  $A$ . This formula had appeared earlier as an exercise in [2, p. 23]. It leads to a simple construction of the mixtilinear incircle. Denote by  $I$  the incenter of triangle  $ABC$ , and let the perpendicular through  $I$  to the bisector of angle  $A$  intersect the sides  $AC$ ,  $AB$  at  $Y_1$  and  $Z_1$ , respectively. The perpendiculars at these points to their respective sides intersect again on the angle bisector, at the mixtilinear incenter  $K_A$ . The circle with center  $K_A$ , passing through  $Y_1$  (and  $Z_1$ ), is the mixtilinear incircle in angle  $A$ ; see Figure 1.

In this note, we demonstrate the usefulness of the notion of barycentric coordinates in discovering remarkable geometric properties relating to the mixtilinear incircles of a triangle. To keep the note self-contained, we refrain from using (1), except for the remarks at the end.

Denote by  $A'$  the point of contact of the mixtilinear incircle in angle  $A$  with the circumcircle. For convenience, we denote  $K_A$  by  $K$ , and  $\rho_A$  by  $\rho$  when there is no danger of confusion; see Figure 2. The center  $K$  lies on the bisector of angle  $A$ , and  $AK : KI = \rho : -(\rho - r)$ . In terms of barycentric coordinates,

$$K = \frac{1}{r} [ -(\rho - r)A + \rho I ]. \tag{2}$$




Figure 1




Figure 2

Also, since the circumcircle  $O(A')$  and the mixtilinear incircle  $K(A')$  touch each other at  $A'$ , we have  $OK : KA' = R - \rho : \rho$ , where  $R$  is the circumradius. From this,

$$K = \frac{1}{R} [\rho O + (R - \rho) A']. \quad (3)$$

Comparing (2) and (3), we obtain, by rearranging terms,

$$\frac{RI - rO}{R - r} = \frac{R(\rho - r)A + r(R - \rho)A'}{\rho(R - r)}. \quad (4)$$

We note some interesting consequences of this formula. First of all, it gives the intersection of the lines joining  $AA'$  and  $OI$ . Note that the point  $P$  on the line  $OI$  represented by the left hand side of (4) is the external center of similitude of the circumcircle and the incircle of the given triangle. This, by definition, is the point dividing the segment  $OI$  externally in the ratio of the radii of the circles. As such, it can be constructed as the intersection of the lines  $OI$  and  $MD$ , where  $M$  is the intersection of the bisector of angle  $A$  with the circumcircle, and  $D$  the point of contact of the incircle with the side  $BC$ ; see Figure 3.

The same reasoning applied to the other two mixtilinear incircles shows that each of the lines  $AA'$ ,  $BB'$ ,  $CC'$  passes through the same point  $P$  on the line  $OI$ ; see Figure 4.




Figure 3




Figure 4

**Theorem 1.** *The three lines each joining a vertex to the point of contact of the circumcircle with the mixtilinear incircle in the angle of the vertex are concurrent at the external center of similitude of the circumcircle and the incircle.*

Equation (4) also leads to an alternative construction of the mixtilinear incircle, without the use of (1).

**Construction 2.** Given a triangle  $ABC$ , let  $P$  be the external center of similitude of the circumcircle ( $O$ ) and incircle ( $I$ ). Extend  $AP$  to intersect the circumcircle at  $A'$ . The intersection of  $AI$  and  $A'O$  is the center  $K_A$  of the mixtilinear incircle in angle  $A$ .

Theorem 1 means that the triangles  $ABC$  and  $A'B'C'$  are in perspective. By Desargues' Theorem, the intersections of the three pairs of lines  $BC, B'C'$ ;  $CA, C'A'$ , and  $AB, A'B'$  are collinear. The intersection  $X$  of the lines  $BC$  and  $B'C'$  is indeed the external center of similitude of the mixtilinear incircles ( $K_B$ ) and ( $K_C$ ). This is clear from the following lemma, whose proof we omit.

**Lemma 3.** *If two distinct circles are tangent to a third circle, both internally or both externally, then the line joining the points of contact passes through the external center of similitude of the two circles.*

If one of the tangencies is internal and the other is external, then the line joining the points of contact passes through the internal center of similitude of the two circles; see Figure 5.

It is easy to determine the barycentric coordinates of  $X$  with respect to  $B$  and  $C$ . In fact,

$$X = \frac{\rho_C \cdot K_B - \rho_B \cdot K_C}{\rho_C - \rho_B} = \frac{-\left(1 - \frac{r}{\rho_B}\right)B + \left(1 - \frac{r}{\rho_C}\right)C}{\left(\frac{1}{\rho_B} - \frac{1}{\rho_C}\right)r}.$$




Figure 5

Here, we have made use of analogues of (2). Similarly, the external centers of similitude of the pairs of circles  $(K_C)$ ,  $(K_A)$ , and  $(K_A)$ ,  $(K_B)$  are

$$Y = \frac{-\left(1 - \frac{r}{\rho_C}\right)C + \left(1 - \frac{r}{\rho_A}\right)A}{\left(\frac{1}{\rho_C} - \frac{1}{\rho_A}\right)r} \quad \text{and} \quad Z = \frac{-\left(1 - \frac{r}{\rho_A}\right)A + \left(1 - \frac{r}{\rho_B}\right)B}{\left(\frac{1}{\rho_A} - \frac{1}{\rho_B}\right)r}.$$

These three points  $X, Y, Z$  all lie on the line

$$\frac{x}{1 - \frac{r}{\rho_A}} + \frac{y}{1 - \frac{r}{\rho_B}} + \frac{z}{1 - \frac{r}{\rho_C}} = 0. \quad (5)$$

Indeed, the triangles  $ABC$ ,  $A'B'C'$ , and  $K_A K_B K_C$  are pairwise in perspective, with line (5) as common axis of perspective.

We close with a few remarks. Since the points  $X, Y, Z$  are the external centers of similitude of pairs of circles from  $(K_A)$ ,  $(K_B)$ ,  $(K_C)$ , their collinearity also follows from the famous Desargues Three-Circle Theorem [5]. If we make use of (1), this axis of perspective has equation

$$\frac{x}{\sin^2 \frac{\alpha}{2}} + \frac{y}{\sin^2 \frac{\beta}{2}} + \frac{z}{\sin^2 \frac{\gamma}{2}} = 0.$$

Finally, we note another interesting consequence of (1). The Gergonne point of a triangle is the point of intersection of the three cevians joining each vertex to the point of contact of the incircle with the opposite side. This is the point  $X_7$  of [4], and has trilinear coordinates

$$\sec^2 \frac{\alpha}{2} : \sec^2 \frac{\beta}{2} : \sec^2 \frac{\gamma}{2}.$$

As such, this is the unique point whose distances to the sides are proportional to the radii of the mixtilinear incircles in the respective angles.

#### REFERENCES

1. L. Bankoff, A Mixtilinear Adventure, *Crux Math.* **9** (1983) 2–7.
2. C. V. Durell and A. Robson, *Advanced Trigonometry*, Bell & Sons, 1935.
3. R. A. Johnson, *Advanced Euclidean Geometry*, Dover reprint, 1965.
4. C. Kimberling, Central Points and Central Lines in the Plane of a Triangle, *Math. Mag.* **67** (1994) 163–187.
5. J. McCleary, An Application of Desargues' Theorem, *Math. Mag.* **55** (1982) 233–235.

Florida Atlantic University, Boca Raton, FL 33431  
 yiu@fau.edu